

CH⁺(1-0) in $z \sim 2-4$ starburst galaxies: probes of multi-phasic turbulent gas reservoirs

Alba Vidal-Garcia¹, Edith Falgarone¹, Fabrizio Arrigoni Battaia², Benjamin Godard¹, Cinthya Herrera³, Martin A. Zwaan⁴, Rob Ivison⁴, Fabian Walter⁵, Edwin Bergin⁶, Paola Andreani⁴, Qiong Li⁷, Alain Omont⁸ and Raphaël Gavazzi⁸

¹ *Laboratoire de Physique de l'Ecole normale supérieure, ENS, Université PSL, CNRS, Sorbonne Université, Université de Paris, F-75005 Paris, France*

² *Max Planck Institute für Astronomie, Garching, Germany*

³ *Institut de radioastronomie millimétrique, 106 rue de la Piscine, Saint Martin d'Hères, France*

⁴ *European Southern Observatory, Karl-Schwarzschild-Strasse 2, 85748 Garching, Germany*

⁵ *Max Planck Institute für Astronomie, Heidelberg, Germany*

⁶ *University of Michigan, Ann Arbor, MI, USA*

⁷ *Kavli Institute for Astronomy and Astrophysics, Peking University, Beijing, People's Republic of China*

⁸ *Sorbonne Université, CNRS, UMR7095, Institut d'Astrophysique de Paris, F-75014, Paris, France*

Starburst galaxies at redshifts $z \sim 2$ to 4 are among the most intensely star-forming galaxies in the universe. The way they accrete their gas to form stars at such high rates is still a controversial issue. We have detected the CH⁺(1-0) line in emission and/or in absorption in all the gravitationally lensed starburst galaxies observed so far with ALMA in this redshift range [1]. The unique spectroscopic and chemical properties of CH⁺ allow its transition to highlight the sites of dissipation of mechanical energy. Whilst the absorption lines reveal highly turbulent reservoirs of low-density molecular gas extending far out of the galaxies, the broad emission lines with widths up to a few thousands of km/s, arise in myriad molecular shocks powered by the feedback of star formation and possibly active galactic nuclei [2,3]. The CH⁺(1-0) lines therefore probe the sites of prodigious energy releases, mainly stored in turbulent reservoirs before being radiated away. These turbulent reservoirs act as extended buffers of mass and energy over timescales of a few tens to hundreds of Myr. Their mass supply involves multiphasic gas inflows from galaxy mergers and/or cold stream accretion, as supported by *Keck*/*KCWI* Ly α observations of one of these starburst galaxies [4,5].

References

- [1] Falgarone, Zwaan et al. 2017, *Nature* 548 430
- [2] Godard, Pineau des Forêts et al. 2019, *A&A* 622, A100
- [3] Lehmann, Godard et al. 2021, *A&A* 658, A165
- [4] Li, Cai, Prochaska et al. 2019, *ApJ* 875 130
- [5] Vidal-García, Falgarone et al. 2021, *MNRAS* 506, Issue 2, 2551-2573